


DIABETES COACH

How to use Bydureon®

Where to inject your weekly dose

Your once-weekly medication is administered as an injection under the skin, just once a week.¹

- You can inject into any of the areas shown in the picture:
 - The stomach area
 - The thigh
 - The back of the upper arm
- You can use the same area of your body every week if you wish. However, you should inject your medicine into a different place within that area each time



Preparing and injecting your once-weekly dose

What's in your single-dose kit


Vial

The vial contains one premeasured dose of powdered Bydureon



Syringe

The liquid will be mixed with the powdered medicine in the vial



Orange vial connector

The vial connector is used to connect the vial with the syringe



Needles (two)

The needle is much shorter than the needle cover; the single-dose kit has a spare needle in case you need it



Connect, Shake and Inject: How the parts fit together


1. Connect the parts securely before mixing



Vial with premeasured dose of medicine



Orange vial connector (and package)



Syringe with liquid (diluent)



Parts connected

2. Shake hard to mix the medicine with the liquid



Shake well



Check that it's mixed well



Not mixed well, keep shaking

3. Inject. Attach the needle, line up the plunger with the dose line and inject subcutaneously

Plunger at black dashed dose line



Remember: Detailed information on how to inject can be found in the *Instructions for Use* booklet that accompanies each single-dose kit.

LOCAL MARKETS TO INSERT WORDING ON ADVERSE-EVENT REPORTING FOR PATIENTS IN ACCORDANCE WITH LOCAL REGULATIONS.

Reference

1. Bydureon®. Summary of product characteristics,
Markets to update with appropriate version upon release.